


Quadratic function

Definition

$f: D \rightarrow \mathbb{R}$	$(D \subseteq \mathbb{R})$
$x \mapsto y = f(x) = ax^2 + bx + c$	$(a \in \mathbb{R} \setminus \{0\}, b \in \mathbb{R}, c \in \mathbb{R})$
general form	
$y = f(x) = a(x - u)^2 + v$	$(a \in \mathbb{R} \setminus \{0\}, u \in \mathbb{R}, v \in \mathbb{R})$
vertex form	

Graph

1. $y = f(x) = x^2$ ($a = 1, u = 0, v = 0$)




2. Parameter **u** (**u is varied**, a and v are kept constant)

$$y = f_0(x) = x^2 \quad (a = 1, u = 0, v = 0)$$

$$y = f_1(x) = (x - 2)^2 \quad (a = 1, u = 2, v = 0)$$

$$y = f_2(x) = (x + 1)^2 \quad (a = 1, u = -1, v = 0)$$




3. Parameter **v** (**v is varied**, a and u are kept constant)

$$y = f_0(x) = x^2 \quad (a = 1, u = 0, v = 0)$$

$$y = f_1(x) = x^2 + 3 \quad (a = 1, u = 0, v = 3)$$

$$y = f_2(x) = x^2 - 2 \quad (a = 1, u = 0, v = -2)$$




4. Parameter **a** (**a is varied**, u and v are kept constant)

$$y = f_0(x) = x^2 \quad (a = 1, u = 0, v = 0)$$


$$y = f_1(x) = 2x^2 \quad (a = 2, u = 0, v = 0)$$

$$y = f_2(x) = -2x^2 \quad (a = -2, u = 0, v = 0)$$



5. Parameter **a** (**a is varied**, u and v are kept constant)

$$\begin{aligned} y = f_0(x) &= x^2 & (a = 1, u = 0, v = 0) \\ y = f_1(x) &= \frac{1}{2}x^2 & (a = \frac{1}{2}, u = 0, v = 0) \\ y = f_2(x) &= -\frac{1}{2}x^2 & (a = -\frac{1}{2}, u = 0, v = 0) \end{aligned}$$




6. The **graph** of a quadratic function is a **parabola**.

The parameter **a** determines the **shape** of the parabola, and whether the parabola opens upwards or downwards.

The parameters **u** and **v** determine the **position** of the parabola. They are the coordinates of the **vertex** V of the parabola: V(u|v)

$$\begin{aligned} y = f_0(x) &= x^2 & (a = 1, u = 0, v = 0) & V(0|0) \\ y = f_1(x) &= 2(x - 2)^2 - 4 & (a = 2, u = 2, v = -4) & V(2|-4) \\ y = f_2(x) &= -\frac{1}{2}(x + 1)^2 + 6 & (a = -\frac{1}{2}, u = -1, v = 6) & V(-1|6) \end{aligned}$$




Examples

1. Nature/Physics: Trajectories of water in a fountain



2. Geometry: Square




Area A at side length l : $A = l^2$

$f: \mathbb{R}^+ \rightarrow \mathbb{R}$

$l \mapsto A = f(l) = l^2$ quadratic function

3. Physics: Free fall



Distance d after time t : $d = \frac{1}{2}gt^2$ (g = gravity field strength)

$f: \mathbb{R} \rightarrow \mathbb{R}$

$t \mapsto d = f(t) = \frac{1}{2}gt^2$ quadratic function

4. Economics: Supply, Demand