

## Exercises 13

### Derivative


#### Rate of change, derivative of constant/power/exponential functions

#### Objectives

- be able to estimate a rate of change out of the graph of a function.
- be able to state the rate of change of a constant/linear function.
- be able to determine a rate of change of a basic power/exponential function.
- be able to determine the derivative of a constant/linear function.
- be able to determine the derivative of a basic power/exponential function.

#### Problems

13.1 The graph of a function  $f$  is given as follows:



Estimate the rate of change  $f'(x_0)$  at the given position  $x_0$ :

- a)  $x_0 = -1$       b)  $x_0 = 0$   
c)  $x_0 = 1$       d)  $x_0 = -2$

Hints:

- Draw the tangent to the graph of  $f$  at the given  $x_0$ .
- Estimate the slope of the tangent.

13.2 The graph of a constant or linear function is a straight line. Therefore, the “tangent” to the graph at any point of the graph is the graph itself.

For each of the following functions  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $x \mapsto y = f(x) = \dots$

- i) ... draw the graph of  $f$ .
  - ii) ... state the rate of change  $f'(x_0)$  at the given  $x_0$ .
- |  |                          |
|--|--------------------------|
| a) $f(x) = 3$  | $x_0 = 2$                |
| b) $f(x) = c$ ( $c \in \mathbb{R}$ )   | any $x_0 \in \mathbb{R}$ |
| c) $f(x) = 2x - 3$   | $x_0 = 4$                |
| d) $f(x) = mx + q$ ( $m \in \mathbb{R} \setminus \{0\}$ , $q \in \mathbb{R}$ ) | any $x_0 \in \mathbb{R}$ |
| e) * $f(x) =  x $  | any $x_0 \in \mathbb{R}$ |

13.3 Determine  $f'(x)$ :

- |                         |                           |                              |
|-------------------------|---------------------------|------------------------------|
| a) $f(x) = 3$           | b) $f(x) = 0$             | c) $f(x) = -1$               |
| d) $f(x) = x^3$         | e) $f(x) = x^4$           | f) $f(x) = x^5$              |
| g) $f(x) = x^{17}$      | h) $f(x) = x^{200}$       | i) $f(x) = x^{100'001}$      |
| j) $f(x) = x^{-1}$      | k) $f(x) = x^{-2}$        | l) $f(x) = x^{-17}$          |
| m) $f(x) = \frac{1}{x}$ | n) $f(x) = \frac{1}{x^3}$ | o) $f(x) = \frac{1}{x^{99}}$ |

13.4 Determine  $f'(x)$ :

- |  |  |  |
|--|--|--|
| a) $f(x) = 3^x$                        | b) $f(x) = 5^x$                          | c) $f(x) = 18^x$                       |
| d) $f(x) = \left(\frac{2}{3}\right)^x$ | e) $f(x) = \left(\frac{13}{17}\right)^x$ | f) $f(x) = \left(\frac{1}{4}\right)^x$ |
| g) $f(x) = \left(\frac{1}{e}\right)^x$ | h) $f(x) = \left(\frac{3}{e}\right)^x$   | i) $f(x) = \left(\frac{e}{3}\right)^x$ |

13.5 Determine the rate of change of the function  $f$  at the indicated  $x_0$ :

- |  |                          |                           |  |  |
|--|--------------------------|---------------------------|--|--|
| a) $f(x) = x$                          |                          |                           |  |  |
| i) $x_0 = 0$                           | ii) $x_0 = 1$            | iii) $x_0 = -2$           |  |  |
| b) $f(x) = x^5$                        |                          |                           |  |  |
| i) $x_0 = 0$                           | ii) $x_0 = 2$            | iii) $x_0 = -\frac{2}{3}$ |  |  |
| c) $f(x) = x^{-4}$                     |                          |                           |  |  |
| i) $x_0 = -1$                          | ii) $x_0 = -\frac{4}{3}$ | iii) $x_0 = 0$            |  |  |
| d) $f(x) = \left(\frac{2}{3}\right)^x$ |                          |                           |  |  |
| i) $x_0 = 0$                           | ii) $x_0 = 1$            | iii) $x_0 = -2$           |  |  |

13.6 \* (see next page)

13.6 \* The rate of change  $f'(x_0)$  of  $f$  at  $x_0$  can be determined by looking at the secant through the points  $A(x_0 | f(x_0))$  and  $B(x_0 + \Delta x | f(x_0 + \Delta x))$  of the graph of  $f$ . The slope of this secant tends towards the slope of the tangent at  $A(x_0 | f(x_0))$  as  $\Delta x$  tends towards 0.

It has been shown in class how to determine  $f'(x_0)$  for the quadratic function  $f(x) = x^2$ .

Find  $f'(x_0)$  for the following functions  $f$ :

a)  $f(x) = x^3$       b)  $f(x) = \frac{1}{x^2}$

13.7 Decide which statements are true or false. Put a mark into the corresponding box.  
In each problem a) to c), exactly one statement is true.

a) The rate of change of the function  $f$  at the position  $x_0$  is a ...

- ... real number.
- ... function.
- ... tangent.
- ... graph.

b) The derivative  $f'$  of the function  $f$  is a ...

- ... real number.
- ... function.
- ... tangent.
- ... graph.

c)  $f'(x_0)$  is the slope of the ...

- ... secant through the points  $(0|0)$  and  $(x_0|f(x_0))$ .
- ... secant through the points  $(x_0 + \Delta x | f(x_0 + \Delta x))$  and  $(x_0 | f(x_0))$ .
- ... tangent to the graph of  $f$  at  $(x_0 | f(x_0))$ .
- ... tangent to the graph of  $f'$  at  $(x_0 | f(x_0))$ .

## Answers

- 13.1      a)       $f'(-1) \approx 0$       b)       $f'(0) \approx 2$   
               c)       $f'(1) \approx \frac{3}{2}$       d)       $f'(-2) \approx -\frac{5}{3}$

- 13.2    a)      i)      ...  
              ii)      $f'(2) = 0$

b)      i)      ...  
              ii)      $f'(x_0) = 0$

c)      i)      ...  
              ii)      $f'(4) = 2$

d)      i)      ...  
              ii)      $f'(x_0) = m$

e) \*     i)      ...  
              ii)      $f'(x_0) = \begin{cases} 1 & (x_0 > 0) \\ -1 & (x_0 < 0) \\ \text{not defined} & (x_0 = 0) \end{cases}$

- 13.3 a)  $f'(x) = 0$       b)  $f'(x) = 0$       c)  $f'(x) = 0$   
d)  $f'(x) = 3x^2$       e)  $f'(x) = 4x^3$       f)  $f'(x) = 5x^4$   
g)  $f'(x) = 17x^{16}$       h)  $f'(x) = 200x^{199}$       i)  $f'(x) = 100'001x^{100'000}$   
j)  $f'(x) = -x^{-2}$       k)  $f'(x) = -2x^{-3}$       l)  $f'(x) = -17x^{-18}$   
m)  $f'(x) = -\frac{1}{x^2}$       n)  $f'(x) = -\frac{3}{x^4}$       o)  $f'(x) = -\frac{99}{x^{100}}$

- 13.4      a)       $f'(x) = 3^x \ln(3)$       b)       $f'(x) = 5^x \ln(5)$       c)       $f'(x) = 18^x \ln(18)$   
d)       $f'(x) = \left(\frac{2}{3}\right)^x \ln\left(\frac{2}{3}\right)$       e)       $f'(x) = \left(\frac{13}{17}\right)^x \ln\left(\frac{13}{17}\right)$   
f)       $f'(x) = \left(\frac{1}{4}\right)^x \ln\left(\frac{1}{4}\right) = -\frac{\ln(4)}{4^x}$

Hint:

- Logarithm rules (see formulary) can be applied in order to simplify the result.

$$g) \quad f'(x) = -\frac{1}{e^x} \qquad h) \quad f'(x) = \left(\frac{3}{e}\right)^x (\ln(3) - 1) \qquad i) \quad f'(x) = \left(\frac{e}{3}\right)^x (1 - \ln(3))$$

- 13.5    a)     $f'(x) = 1$

i)     $f'(0) = 1$               ii)     $f'(1) = 1$               iii)     $f'(-2) = 1$

b)     $f'(x) = 5x^4$

i)     $f'(0) = 0$               ii)     $f'(2) = 80$               iii)     $f'\left(-\frac{2}{3}\right) = \frac{80}{81}$

c)     $f'(x) = -\frac{4}{x^5}$

i)     $f'(-1) = 4$               ii)     $f'\left(-\frac{4}{3}\right) = \frac{243}{256}$               iii)     $f'(0)$  is not defined

d)    (see next page)

d)  $f'(x) = \left(\frac{2}{3}\right)^x \ln\left(\frac{2}{3}\right)$

i)  $f'(0) = \ln\left(\frac{2}{3}\right)$       ii)  $f'(1) = \frac{2}{3} \ln\left(\frac{2}{3}\right)$       iii)  $f'(-2) = \frac{9}{4} \ln\left(\frac{2}{3}\right)$

13.6 \* a) ... b) ...

- 13.7      a)      1<sup>st</sup> statement  
              b)      2<sup>nd</sup> statement  
              c)      3<sup>rd</sup> statement