


Quadratic function

Definition

$f: D \rightarrow \mathbb{R}$	$(D \subseteq \mathbb{R})$
$x \mapsto y = f(x) = ax^2 + bx + c$	$(a \in \mathbb{R} \setminus \{0\}, b \in \mathbb{R}, c \in \mathbb{R})$
	general form
$y = f(x) = a(x - u)^2 + v$	$(a \in \mathbb{R} \setminus \{0\}, u \in \mathbb{R}, v \in \mathbb{R})$
	vertex form


Graph

1. $y = f(x) = x^2$ ($a = 1, u = 0, v = 0$)




2. Parameter u

$y = f_0(x) = x^2$	$(a = 1, u = 0, v = 0)$
$y = f_1(x) = (x - 2)^2$	$(a = 1, u = 2, v = 0)$
$y = f_2(x) = (x + 1)^2$	$(a = 1, u = -1, v = 0)$




3. Parameter **v**

$$\begin{aligned}y &= f_0(x) = x^2 & (a = 1, u = 0, v = 0) \\y &= f_1(x) = x^2 + 3 & (a = 1, u = 0, v = 3) \\y &= f_2(x) = x^2 - 2 & (a = 1, u = 0, v = -2)\end{aligned}$$




4. Parameter **a**

$$\begin{aligned}y &= f_0(x) = x^2 & (a = 1, u = 0, v = 0) \\y &= f_1(x) = 2x^2 & (a = 2, u = 0, v = 0) \\y &= f_2(x) = -2x^2 & (a = -2, u = 0, v = 0)\end{aligned}$$



5. Parameter **a**

$$\begin{array}{ll} y = f_0(x) = x^2 & (\mathbf{a = 1}, u = 0, v = 0) \\ y = f_1(x) = \frac{1}{2}x^2 & (\mathbf{a = \frac{1}{2}}, u = 0, v = 0) \\ y = f_2(x) = -\frac{1}{2}x^2 & (\mathbf{a = -\frac{1}{2}}, u = 0, v = 0) \end{array}$$




6. The **graph** of a quadratic function is a **parabola**.

The parameter **a** determines the **shape** of the parabola, and whether the parabola opens upwards or downwards.

The parameters **u** and **v** determine the **position** of the parabola. They are the coordinates of the **vertex** V of the parabola: $V(u|v)$

$$\begin{array}{lll} y = f_0(x) = x^2 & (\mathbf{a = 1}, u = 0, v = 0) & V(0|0) \\ y = f_1(x) = 2(x - 2)^2 - 4 & (\mathbf{a = 2}, u = 2, v = -4) & V(2|-4) \\ y = f_2(x) = -\frac{1}{2}(x + 1)^2 + 6 & (\mathbf{a = -\frac{1}{2}}, u = -1, v = 6) & V(-1|6) \end{array}$$




Examples

1. Nature/Physics: Trajectories of water in a fountain




2. Geometry: Square



Area A for side length l : $A = l^2$

$$f: \mathbb{R}^+ \rightarrow \mathbb{R}$$
$$l \mapsto A = f(l) = l^2 \quad \text{quadratic function}$$

3. Physics: Free fall



Distance d after time t : $d = \frac{1}{2}gt^2$ (g = gravity field strength)

$$f: \mathbb{R} \rightarrow \mathbb{R}$$
$$t \mapsto d = f(t) = \frac{1}{2}gt^2 \quad \text{quadratic function}$$

4. Economics: Supply, Demand